

Lighting And Material Of Halo3

Hao Chen (haochen@bungie.com)

Bungie

Microsoft Research Asia



NON FACETE NOBIS CALCITRARE VESTRUM

Talk Overview

- Introduction
- Halo3 Lighting
- Halo3 Material Model
- HDR Rendering
- Results
- Acknowledgement
- Q&A

BUNGIE

NON FACETE NOBIS CALCITRARE VESTRUM



Halo3

- What do we want?
 - global illumination
 - handle variety of lighting environments
 - consistent lighting everywhere
 - render bump maps “correctly”
 - complex material under complex lighting
 - HDR

BUNGIE

NON FACETE NOBIS CALCITRARE VESTRUM

Definitions

- Radiance
 - $L(\omega)$: density of light energy through a given point in a given direction.
- Irradiance
 - I : incident radiance integrated over the hemisphere of the surface normal with the cosine lobe.
- BRDF
 - $f(V, L)$: Bidirectional reflectance distribution function.
- Fresnel
 - F : Predicts ratio of reflected and transmitted light when light travels between different mediums.
 - $F0$: Reflectance at near normal incident angle.

Related Work

- Irradiance Volume [Greger98][ATI05]
- SH Irradiance Environment Map [Ramamoorthi01]
- Pre-computed Radiance Transfer [Sloan02]
- SH Light Maps [Good, Taylor05]
- Sky and Atmosphere [Preetham99][Hoffman02]
- Reflectance Models [CookTorrance82][Schlick94]
- Low Frequency Glossy Material [Kautz02][Sloan03]
- Frequency Space Environment Map [Ramamoorthi02]

SH Irradiance Env Map



[Ramamoorthi00]

$$L_{lm} = \int_{\theta, \phi} L(\theta, \phi) Y_{lm}(\theta, \phi) \sin(\theta) d\theta d\phi$$

distant radiance $L(\theta, \phi)$ at given direction (θ, ϕ) $Y_{lm}(\theta, \phi)$ $\sin(\theta) d\theta d\phi$ diff solid angle

SH Irradiance Env Map




$$E(n) = n^T M n$$

- Irradiance distribution as SH vector.
- <3% error with just 9 terms [Basri Jacobs 01]
- Evaluate normal to get irradiance.
- Only represent a single point in space.
- Only for infinite lighting environment.
- What about local lighting?

Irradiance Volumes



[Greger 98]




[ATI 05]

- Spatially divide volume into cells.
- irradiance volume per cell.
- Interpolate between samples.
- Sharp shadow boundaries?
- Bump maps?



SH Light Map



Parameterize



GI Solver



GI Solver




GI Solver




GI Solver



Rendering



Compression



GI Solver



GI Solver



GI Solver


Other basis would work too, e.g. Half Life Basis, ZH, etc.

BUNGIE

NON FACETE NOBIS CALCITRARE VESTRUM

Sky and Sun

- HDR pipeline means starting with HDR light sources.
- Custom sky plugin for 3DStudio Max.
- Procedural Sky/Sun Model[Preetham99]
- CIE Sky Model
- Can also use HDR Cubemap




HDR Sky Example



BUNGIE

NON FACETE NOBIS CALCITRARE VESTRUM

Parameterization



- UVAtlas (DirectX SDK)

- Minimize distortion
- Efficient Packing
- Input "importance"

- Halo3 improvements:

- small charts placement.
- long and thin charts.


- > 80% texel utilization

- Halo2: < 50%

BUNGIE

NON FACETE NOBIS CALCITRARE VESTRUM

Photon Mapping Farm




256 servers; 456 Processors; 1GB memory per proc; see Luis's talk.

BUNGIE

NON FACETE NOBIS CALCITRARE VESTRUM

Compression



Signal Optimization:


- Compute signal gradient
- Resample to half dim.
- Preserve high freq charts
- Squeeze low freq charts

DXT Compression

- Use 2 DXT5 for each FP16
- Color space conversion
- 12 bits (2 DXT5 alpha) luminance
- 3:1 compression ratio

Details in a separate talk by Yaohua Hu.


Rendering Passes



BUNGIE

NON FACETE NOBIS CALCITRARE VESTRUM

Object Lighting



Optimization

- given a quadratic SH vecotor, $i=0, \dots, 8$
 - Pull out dominant light.
 - Store SH linear + dominant light.
 - In Shader, do $N \cdot L + \text{shirm}[\text{sh}[] - c * Y(d), N]$

$$E = \sum_{i=0, \dots, 8} (\lambda_i - c Y_i(d))^2, E' = 0$$

$$c = \sum_{i=0, \dots, 8} (\lambda_i Y_i(d)) / \sum_{i=0, \dots, 8} Y_i(d)^2$$



See Peter Pike Sloan's talk "Stupid Spherical Harmonics Tricks".

BUNGIE

NON FACETE NOBIS CALCITRARE VESTRUM

Lighting Examples



BUNGIE

NON FACETE NOBIS CALCITRARE VESTRUM

Lighting Examples



BUNGIE

NON FACETE NOBIS CALCITRARE VESTRUM

Lighting Examples



BUNGIE

NON FACETE NOBIS CALCITRARE VESTRUM


Lighting Examples



BUNGIE

NON FACETE NOBIS CALCITRARE VESTRUM


Lighting Examples



BUNGIE

NON FACETE NOBIS CALCITRARE VESTRUM


Lighting Examples



BUNGIE

NON FACETE NOBIS CALCITRARE VESTRUM

Material Motivation



Halo3 Material Model

Diffuse



Area Specular



Environment Map



Analytical Specular



- The basic Idea

- Separate material into diffuse parts and
- Low, med, high freq glossy parts.
- SH irradiance envmap for diffuse
- New area specular model for low frequency glossy.
- Prefiltered env map for mid frequency glossy.
- BRDF evaluated directly with point lights for high freq.


BUNGIE

NON FACETE NOBIS CALCITRARE VESTRUM

Cook Torrance BRDF



[CookTorrance81]



[Ngan05]

$$f(V, L) = k_d R_d + k_s \cdot F \cdot R_m$$

view direction V light direction L k_d R_d k_s F R_m

$$R_m(V, L) = \frac{D G}{\pi (N \cdot L)(N \cdot V)}$$

D: microfacet distribution function

G: geometry term

Cook Torrance & Point Lights

- Option1: Evaluate directly in shader
 - E.g. [Dempski Viale 2005]
 - Somewhat expensive, not too bad.
- Option2: Store D, G, F, terms in textures.
- What about lights that are not point lights?
 - Need to integrate light from all directions.
 - Not trivial to do.

Light Integration

$$I(V) = \oint f(V, L) \cos(\theta) L(\omega) d\omega$$

$$k_d R_o \int f(V, L) \cos(\theta) L(\omega) d\omega + k_s \int F(R, V, L) \cos(\theta) L(\omega) d\omega$$



??



SH irradiance env. map

BUNGIE

NON FACETE NOBIS CALCITRARE VESTRUM

Light Integration in SH

$$I_s(V) = k_s \oint FR_m(V, L) \cos(\theta) L(\omega) d\omega$$

$$L(\omega) = \sum_{i=0}^8 \lambda_i Y_i(\omega)$$

Project light into SH basis.

$$B_{m,i}(V) = \oint \frac{F}{F_0} R_m(V, L) \cos(\theta) Y_i(\omega) d\omega$$

Project BRDF and cosine term in SH basis

$$I_s(V) = K_s F_0 \sum_{i=0}^8 \lambda_i B_{m,i}(V) \leftarrow \text{Dot product to convolve}$$

Light Integration in SH Cont.

$$F \approx F_0 + (1 - F_0)(1 - (L \cdot H))^5 \quad [\text{Shlick85}]$$

$$B_{m,i}(V) = \int \left[F_0 + (1 - F_0)(1 - (L \cdot H))^5 \right] R_m(V, L) \cos(\theta) Y_i(\omega) d\omega$$

$$C_{m,i}(V) = \int R_m(V, L) \cos(\theta) Y_i(\omega) d\omega$$

$$D_{m,i}(V) = \int (1 - (L \cdot H)^5) R_m(V, L) \cos(\theta) Y_i(\omega) d\omega$$

$$B_{m,i}(V) = C_{m,i}(V) + \frac{1 - F_0}{F_0} D_{m,i}(V) \quad \leftarrow \text{Preintegration}$$

Pre-integration



Isotropic BRDF = any coordinate frame

Reflective symmetry means:

$$C_{m,i}(V) = D_{m,i}(V) = 0, i = 1, 4, 5.$$


16 m values, and 8 V directions is enough.



C (i=0,2,3,6)




D (i=0,2,3,6)




C,D (i=7,8)


BUNGIE

NON FACETE NOBIS CALCITRARE VESTRUM














Rendering Pipeline



Render Target Considerations

- Memory Size
 - Render Speed
 - Hardware Blend Support
 - Dynamic Range
 - Banding
- } Useable Exposure Range

Xbox 360 Render Targets

	16f	10-bit 7e3	10-bit lin.	8-bit xRGB	16-bit lin.
Exposure Range	30 stops	3 stops	0 stops	0 stops	5 stops
Blend Support	NO ☹	YES	YES	YES	YES
Memory Size	2x	1x	1x	1x	2x
Blend/Fill Rate	50%	50-100%	50-100%	50-100%	50%

	8-bit xRGB + 8-bit xRGB
Exposure Range	7 stops
Blend Support	YES
Memory Size	2x
Blend/Fill Rate	25-50%




BUNGIE


NON FACETE NOBIS CALCITRARE VESTRUM








Bloom Postprocessing



BUNGIE

NON FACETE NOBIS CALCITRARE VESTRUM

















Conclusion

- SH light map is a natural extension to the traditional light-mapping pipeline.
- Separating material into layers is a good approximation for all frequency reflectance.
- Area specular is critical for achieving seamless lighting and material integration.
- ALU is cheap, and will get cheaper, take maximum advantage of it.

Future Work

- Global Illumination with local, moving lights.
- GI for dynamic and semi-dynamic scenes.
- Better lighting basis (less ringing, higher frequency).
- Area specular model with complex transport.
- Measured BRDF.
- Non photo-realistic rendering.

Acknowledgement

- Authors

- **Bungie**: Hao Chen, Ben Wallace, Chris Tchou, David Cook, Xi Wang, Yaohua Hu.
- **MSRA**: Xinguo Liu, Zhipeng Hu, Xin Huang, Minmin Gong, Kun Zhou.

- Special Thanks

- Bungie artists.
- Peter Pike Sloan, Baining Guo, Harry Shum
- Kutta Srinivasan, Matt Lee, Mikey Wetzel.



These slides will be available online
Feb. 25th, 2008

<http://www.bungie.net/publications/>

We're Hiring!

- Come see us at the Career Pavillion in
Booth 913 (West Hall)

<http://www.bungie.net/jobs/>



NON FACETE NOBIS CALCITRARE VESTRUM

Q & A

BUNGIE

NON FACETE NOBIS CALCITRARE VESTRUM